

TRANSMITTAL LETTER TO THE UNITED STATES

DESIGNATED/ELECTED OFFICE (DO/EO/US)

CONCERNING A FILING UNDER 35 U.S.C. 371

61765.00385

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR

10/030414

INTERNATIONAL APPLICATION NO.

PCT/US00/18817

INTERNATIONAL FILING DATE

July 11, 2000

PRIORITY DATE CLAIMED

July 12, 1999

TITLE OF INVENTION

2-AMINO-BENZOXAZINONES FOR THE TREATMENT OF HERPES SIMPLEX VIRUS

APPLICANT(S) FOR DO/EO/US

Masashi KAWANISHI and Wataru TAKAHASHI

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (24) indicated below.
4. ☒ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371 (c) (2))
 - a. ☒ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☒ has been communicated by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☐ is attached hereto.
 - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).
11. ☐ A copy of the International Preliminary Examination Report (PCT/IPEA/409).
12. ☒ A copy of the International Search Report (PCT/ISA/210).

Items 13 to 20 below concern document(s) or information included:

13. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
14. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
15. ☒ A **FIRST** preliminary amendment.
16. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
17. ☐ A substitute specification.
18. ☐ A change of power of attorney and/or address letter.
19. ☐ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
20. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
21. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
22. ☐ Certificate of Mailing by Express Mail
23. ☒ Other items or information:

PCT/RO/101 (5 pp.); PCT/RO/401 (5 pp.); PCT/IPEA/436 (1 p.); PCT/IB/306 (1 p.); PCT/IB/304 (1 p.); PCT/IB/308 (1 p.); PCT/IOPEA/402 (1 p.); PCT/IB/301 (2 pp.); Copy of WO 01/03697 published January 18, 2001 w/ISR: Specification (40 pp.), Claims 1-9 (6 pp.), Abstract (1 p.)

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR

INTERNATIONAL APPLICATION NO.

ATTORNEY'S DOCKET NUMBER

10/030414

PCT/US00/18817

61765.00385

24. The following fees are submitted:

CALCULATIONS PTO USE ONLY

BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) :

- ☐ Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1040.00
- ☐ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$890.00
- ☒ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$740.00
- ☐ International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$710.00
- ☐ International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00

ENTER APPROPRIATE BASIC FEE AMOUNT =**\$740.00**

Surcharge of **\$130.00** for furnishing the oath or declaration later than ☐ 20 ☐ 30 months from the earliest claimed priority date (37 CFR 1.492 (e)).

\$0.00

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	9 - 20 =	0	x \$18.00
Independent claims	2 - 3 =	0	x \$84.00
Multiple Dependent Claims (check if applicable).			<input type="checkbox"/>

\$0.00**TOTAL OF ABOVE CALCULATIONS =****\$740.00**

- ☐ Applicant claims small entity status. See 37 CFR 1.27). The fees indicated above are reduced by 1/2.

\$0.00**SUBTOTAL =****\$740.00**

Processing fee of **\$130.00** for furnishing the English translation later than ☐ 20 ☐ 30 months from the earliest claimed priority date (37 CFR 1.492 (f)).

\$0.00**TOTAL NATIONAL FEE =****\$740.00**

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable).

\$0.00**TOTAL FEES ENCLOSED =****\$740.00**

Amount to be:
refunded \$
charged \$

- a. ☐ A check in the amount of _____ to cover the above fees is enclosed.
- b. ☒ Please charge my Deposit Account No. 19-0733 in the amount of \$740.00 to cover the above fees. A duplicate copy of this sheet is enclosed.
- c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 19-0733. A duplicate copy of this sheet is enclosed.
- d. ☐ Fees are to be charged to a credit card. **WARNING:** Information on this form may become public. **Credit card information should not be included on this form.** Provide credit card information and authorization on PTO-2038.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

BANNER & WITCOFF, LTD.
1001 G Street, N.W., 11th Floor
Washington, D.C. 20001

SIGNATURE

Susan A. Wolffe

NAME

33,568

REGISTRATION NUMBER

January 10, 2002

DATE

Attorney Docket No. 61765.00385
International Application No. PCT/US00/18817

PATENT APPLICATION

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of

BOX PCT

Masashi KAWANISHI, et al.

**National Phase Application
PCT/US00/18817
Filed: July 11, 2000**

Serial No.: Unassigned

Group Art Unit: Unassigned

Filed: CONCURRENTLY HEREWITH

Examiner: Unassigned

For: **2-AMINO-BENZOXAZINONES FOR THE TREATMENT OF HERPES SIMPLEX
VIRUS**

PRELIMINARY AMENDMENT

Assistant Honorable Commissioner of
Patents and Trademarks
Washington, D.C. 20231

Sir:

Preliminarily to the examination of the above-identified application, kindly amend the
application as follows:

In the Specification:

Page 1, after the title, insert the following paragraph:


--This is a U.S. National Phase Application Under 35 USC 371 and applicants herewith
claim the benefit of priority of PCT/US00/18817 filed July 11, 2000, which was published Under
PCT Article 21(2) in English and Application No. 60/142,956 filed in the United States of America
on July 12, 1999.--

REMARKS

The amendment to the specification is made in accordance with 35 U.S.C. 119, 37 C.F.R.

1.78 and 37 C.F.R. 1.55. Entry is requested.

Respectfully submitted,


Susan A. Wolfe
Reg. No. 33,568

January 10, 2002
BANNER & WITCOFF, LTD.
Eleventh Floor
1001 G Street, N.W.
Washington, D.C. 20001-4597
(202) 508-9100

**2-AMINO-BENZOXAZINONES
FOR THE TREATMENT OF HERPES SIMPLEX VIRUS**

Field of the Invention

This invention is in the field of antiviral agents and specifically relates to compounds, compositions and methods for treating Herpes Simplex Virus.

Background of the Invention

There is a great need for new therapies for the treatment of viral diseases. Whereas there has been great progress in developing a variety of therapies for the treatment of bacterial infections, there are few viable therapies for the treatment of viruses. Zidovudine is the primary approved treatment for human immunodeficiency virus. Ganciclovir, acyclovir, and foscarnet are currently utilized for the treatment of herpesvirus infections. However, these therapies can have substantial side effects based on their deleterious effects on host cell DNA replication or their effect on a limited number of viral infections. In addition, viruses are known to develop resistance to therapies, which causes a progressive decline in efficacy.

Viruses are classified into broad categories based on whether they incorporate RNA or DNA. Important virus families classified of the DNA type include adenoviridae, poxviridae, papovaviridae and herpesviridae.

Herpesviridae is a family of DNA viruses which include herpes simplex virus type-1 (HSV-1), herpes simplex virus type-2 (HSV-2), cytomegalovirus (CMV), varicella-zoster virus (VZV), Epstein-Barr virus, human herpesvirus-6 (HHV6), human herpesvirus-7 (HHV7), human herpesvirus-8 (HHV8), pseudorabies and rhinotracheitis, among others.

It is known that herpesviruses express their genetic content by directing the synthesis of a number of proteins encoded by the herpesvirus DNA in the host cell. One of the important virus-encoded proteins is made as a precursor consisting of an amino terminal-located protease and carboxyl terminal-located assembly protein. This precursor is proteolytically processed in an autocatalytic manner at a specific

amino acid sequence known as the "release" site yielding separate protease and assembly protein. The assembly protein is cleaved further by the protease at another specific amino acid sequence known as the "maturation" cleavage site. Recently, EP 514,830, published November 25, 1992, describes a virus-specific serine protease which has a role in herpesvirus replication. Additionally, Lui and Roizman (*J. Virol*, 65, 5149 (1991)) describe the sequence and activity of a protease and the associated assembly protein encoded by U_L26 of HSV-1. A. R. Welch et al. (*Proc. Natl. Acad. Sci. USA*, 88, 10792 (1991) and WO93/01291, published January 21, 1993) describe the related protease (also known as assemblin) and assembly protein encoded by U_L80 of CMV. An approach currently being investigated for potential use in the treatment of herpesvirus infections is the development of inhibitors of herpesvirus proteases.

4H-3,1-Benzoxazinones have been described in the literature as having serine protease activity, among others. For example, Teshima et al. (*J. Biol. Chem.*, 257, 5085-5091 (1982)) describes various 2-alkyl-4H-3,1-benzoxazin-4-ones as enzyme inhibitors. Moorman and Abeles (*J. Amer. Chem. Soc.*, 104, 6785-6786 (1982)) describes 4H-3,1-benzoxazin-2,4-dione as having some enzyme inhibitory activity. R. Stein, et al. (*Biochemistry*, 26, 4126-4130, (1987)) describes 2-alkyl-4H-benzoxazin-4-ones, with further substitution at the 5, 6 and 7 positions, as inhibiting the elastase enzyme. WO 92/18488 (published October 29, 1992) describes 2-alkyl-4H-3,1-benzoxazin-4-ones with substitution at the 5 and 7 positions as selective inhibitors of elastase. EP Pub. 206,323 (published December 30, 1985) describes 2-alkoxy-, 2-aryloxy- and 2-aralkoxy-4H-3,1-benzoxazin-4-ones, having substitution at the 5, 6, 7 and 8 positions, as enzyme inhibitors. USP 4,745,116 to A. Kranz et al. describes 2-alkoxy, 2-aryloxy- and 2-aralkoxy-4H-3,1-benzoxazin-4-ones, having further substitution at the 5, 7 and 8 positions, as enzyme inhibitors. USP 5,428,021 to Hiebert et al. describes 6-(aminoacid)amino-2-alkoxybenzoxazinones as elastase inhibitors. WO 96/07648, published March 14, 1996, describes 2-phenylamino-benzoxazinones for the treatment of Alzheimer's, and specifically 6-chloro-2-(2-iodophenylamino)-benzo[d][1,3]oxazin-4-one is described.

2-Amino-4*H*-3,1-benzoxazinones have been described. A. Krantz et al. (*J. Med. Chem.*, 33, 464-479 (1990)) describes 4*H*-3,1-benzoxazin-4-ones substituted with alkyl, alkylamino, alkoxy and alkylthio substituents at the 2-position, and with further substitution at the 5, 6 and 7 positions, as elastase inhibitors. Uejima et al. (*J. Pharm. Exp. Ther.*, 265, 516-522 (1993)) describe 2-alkylamino-5-methyl-7-acylamino-4*H*-3,1-benzoxazin-4-ones as highly selective elastase inhibitors with significant plasma stability. USP 4,657,893 to Krantz et al. describes 2-alkylamino- and 2-alkylurido-4*H*-3,1-benzoxazin-4-ones having further substitution at the 5, 7 and 8 positions, as enzyme inhibitors.

F. L. M. Alvarez (*An. Quim.*, 79, 115-17 (1983)) describes the preparation of 2-sulfonylamino-4*H*-3,1-benzoxazinones. J. G. Tercero et al. (*An. Quim.*, 83, 247-50, (1987)) describes the preparation of 2-arylsulfonylamino-4*H*-3,1-benzoxazinones.

I. Butula et al. (*Croat. Chem. Acta*, 54, 105-8 (1981)) describe the synthesis of 2-alkylamino-4*H*-3,1-benzoxazinones. H. Urich et al. (*J. Org. Chem.*, 32, 4052-53 (1967)) describes the synthesis of 2-alkylamino-4*H*-3,1-benzoxazinones. E. Papadopoulos (*J. Heterocyclic. Chem.*, 21, 1411-14 (1984)) describes the use of 2-haloalkylamino-4*H*-3,1-benzoxazin-4-one as a starting material for the synthesis of phenylureas. EP Appln. 466,944 (published January 22, 1992) describes 2-alkylamino-7-acylamino-5-alkyl-4*H*-benzoxazin-4-ones as selective enzyme inhibitors of elastase.


M. Badawy et al. (*J. Heterocyclic. Chem.*, 21, 1403-4 (1984)) describe the use of N-phenyl-2-amino-4*H*-3,1-benzoxazin-4-one as a starting material for the synthesis of quinazolines. R. Khan et al. (*J. Chem. Research(S)*, 342-43 (1992)) describe the synthesis of 2-[5-aryl-1,3,4-oxadiazol-2-yl]amino-4*H*-3,1-benzoxazin-4-ones.

WO 96/37485 describes antiviral agents and compounds, compositions, and methods for treating herpes-related disorders. This document does not describe compounds that have specificity to HSV.

Summary of the Invention

It was discovered that specific compounds that fall within the generic scope of Formula II of WO 96/37485 are very effective against HSV at lower concentrations than the compounds of the examples specifically disclosed in WO 96/37485. The compounds of the present invention have specificity to HSV. This specificity is deemed to be derived from structural conformation of the compounds having R²⁸ and R²⁹ substituents.


The present invention is directed to a compound of Formula II:



II

wherein R²⁸ is selected from amino optionally substituted with two radicals selected from alkyl, aralkyl, heterocyclalkyl, heterocycl, and aryl;

wherein R²⁹ is selected from




wherein R³⁰ is selected from alkyl, alkoxy, alkylamino, carboxyalkyl, alkoxyalkyl, alkylaminoalkyl, cycloalkyl, heterocycl, heterocyclalkyl, aralkyl, aralkoxy, aryloxy, cycloalkyloxy, arylamino, aralkenyl, heterocyclalkoxy, alkylaminoalkoxy, alkylaminoalkylamino, heterocyclalkylamino, N-aryl-N-alkylamino, and N-aralkylamino; wherein R³¹ is alkyl; wherein R³² is selected from alkyl and aryl; and wherein R³³ is selected from hydrido, halo and alkyl; or a pharmaceutically-acceptable salt thereof.

The present invention is further directed to a pharmaceutical composition comprising a therapeutically-effective amount of a compound of formula II and a pharmaceutically acceptable carrier or diluent.

The present invention is further directed to a method or therapeutic or prophylactic treatment of Herpes Simplex Virus in a subject, said method comprising treating said subject with an effective amount of a compound of formula II.

Detailed Description of the Invention


The present invention relates to a class of substituted benzoxazinones, useful in the therapeutic and prophylactic treatment of Herpes Simplex Virus viral infections, as defined by Formula II:



II

wherein R^{28} is selected from amino optionally substituted with two radicals selected from alkyl, aralkyl, heterocyclalkyl, heterocycl, and aryl, or the nitrogen can form a member of a heterocyclic ring;


wherein R^{29} is selected from



wherein R^{30} is selected from alkyl, alkoxy, alkylamino, carboxyalkyl, alkoxyalkyl, alkylaminoalkyl, cycloalkyl, heterocycl, heterocyclalkyl, aralkyl, aralkoxy, aryloxy, cycloalkyloxy, arylamino, aralkenyl, heterocyclalkoxy, alkylaminoalkoxy, alkylaminoalkylamino, heterocyclalkylamino, N-aryl-N-

alkylamino, and N-aralkylamino; wherein R^{31} is alkyl; wherein R^{32} is selected from alkyl and aryl; and wherein R^{33} is selected from hydrido, halo and alkyl; or a pharmaceutically-acceptable salt thereof.

An even more preferred class of compounds consists of those compounds of Formula II wherein R^{28} is selected from amino optionally substituted with two radicals selected from lower alkyl, lower aralkyl, lower heterocyclalkyl, heterocycl, and aryl, wherein R^{29} is selected from



wherein R^{30} is selected from lower alkyl, lower alkoxy, lower alkylamino, lower carboxyalkyl, lower alkoxyalkyl, lower alkylaminoalkyl, lower cycloalkyl, heterocycl, lower heterocyclalkyl, lower heterocyclalkoxy, lower aralkenyl, lower aralkyl, lower aralkoxy, phenyloxy, phenylamino, lower cycloalkyloxy, lower N-phenyl-N-alkylamino, lower alkylaminoalkoxy, lower alkylaminoalkylamino, lower heterocyclalkylamino, and lower N-aralkylamino; wherein R^{31} is lower alkyl; wherein R^{32} is selected from lower alkyl and aryl; and

wherein R^{33} is selected from hydrido and lower alkyl; or a pharmaceutically-acceptable salt thereof.

Most preferred are compounds where R^{28} is selected from the group consisting of methyl(phenylmethyl)amino, methyl[(4-methoxyphenyl)-methyl]amino, 1-(1,2,3,6-tetrahydro-pyridyl), isopropyl(methyl)amino, (4-furoyl)piperazinyl, (2-cyano)ethyl, (4-thenoyl)piperazinyl, (4-benzenesulfonyl)-piperazinyl, diisopropylamino, [methyl(4-dimethylamino)phenylmethyl]amino, methyl(2-pyridylmethyl)-amino; methyl[2-(3-indolyl)ethyl]amino, 4-morpholyl, allyl(methyl)amino, 1-decahydroquinolyl, and 4-(1-acetyl piperadiny); and R^{29} is selected from the group consisting of [(1,1-dimethylethoxy)carbonyl]amino, benzoylamino, (phenylmethoxyacetyl)amino, and (2,4,6-Trifluorobenzoyl)amino.

The term "hydrido" denotes a single hydrogen atom (H). This hydrido radical may be attached, for example, to an oxygen atom to form a hydroxyl radical or two hydrido radicals may be attached to a carbon atom to form a methylene (-CH₂-) radical. Where used, either alone or within other terms such as "haloalkyl", "alkylsulfonyl", "alkoxyalkyl", "hydroxyalkyl" and "aralkyl" the term "alkyl" embraces linear or branched radicals having one to about twenty carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about ten carbon atoms. Most Preferred are lower alkyl radicals having one to about six carbon atoms. Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, *tert*-butyl, pentyl, iso-amyl, hexyl and the like.

The term "halo" means halogens such as fluorine, chlorine, bromine, or iodine.

The terms "alkoxy" and "alkoxyalkyl" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms. More preferred alkoxy radicals are "lower alkoxy" radicals having one to six carbon atoms. Examples of such radicals include methoxy, ethoxy, propoxy, butoxy and *tert*-butoxy.

The term "alkoxyalkyl" also embraces alkyl radicals having two or more alkoxy radicals attached to the alkyl radical, that is, to form monoalkoxyalkyl and dialkoxyalkyl radicals. More preferred alkoxyalkyl radicals are "lower alkoxyalkyl" radicals having one to six carbon atoms and one or two alkoxy radicals. Examples of such radicals include methoxymethyl, methoxyethyl, ethoxyethyl, methoxybutyl and methoxypropyl. The "alkoxy" or "alkoxyalkyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide "haloalkoxy" or haloalkoxyalkyl radicals. More preferred haloalkoxy radicals are "lower haloalkoxy" radicals having one to six carbon atoms and one or more halo radicals. Examples of such radicals include fluoromethoxy, chloromethoxy, trifluoromethoxy, trifluoroethoxy, fluoroethoxy and fluoropropoxy.

The term "aryl", alone or in combination, means a carbocyclic aromatic system containing one, two or three rings wherein such rings may be attached together in a pendent manner or may be fused.

The term "aryl" embraces aromatic radicals such as phenyl, naphthyl, tetrahydronaphthyl, indane and biphenyl. Aryl moieties may also be substituted at a substitutable position with one or more substituents selected independently from alkyl, aralkyl, alkoxyalkyl, alkylaminoalkyl, carboxyalkyl, alkoxycarbonylalkyl, aminocarbonylalkyl, alkoxy, aralkoxy, heterocyclalkoxy, alkylaminoalkoxy, carboxyamino, carboxyaminalkyl, carboxyaminaralkyl, amino, halo, nitro, alkylamino, alkylcarbonylamino, arylcarbonylamino, alkoxycarbonylamino, aralkoxycarbonylamino, aminocarbonylamino, alkylaminocarbonylamino, alkylsulfonylamino, arylsulfonylamino, acyl, cyano, carboxy, aminocarbonyl, alkylaminocarbonyl, alkoxycarbonyl and aralkoxycarbonyl.

The terms "heterocycl" or "heterocyclic" embrace saturated, partially saturated and unsaturated heteroatom-containing ring-shaped radicals, where the heteroatoms may be selected from nitrogen, sulfur and oxygen. Examples of saturated heterocyclic radicals include saturated 5 to 7-membered heteromonocyclic group containing 1 to 4 nitrogen atoms (e.g., pyrrolidinyl, imidazolidinyl, piperidinyl, piperazinyl, tropanyl, homotropanyl, etc.); saturated 5 to 7-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms (e.g., morpholinyl, etc.); saturated 5 to 7-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms (e.g., thiazolidinyl, etc.). Examples of partially saturated heterocyclic radicals include dihydrothiophene, dihydropyran, oxazoliny, dihydrofuran and dihydrothiazole. Examples of unsaturated heterocyclic radicals, also termed "heteroaryl" radicals include unsaturated 5 to 7-membered heteromonocyclic group containing 1 to 4 nitrogen atoms, for example, pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, pyrimidyl, azepinyl, pyrazinyl, pyridazinyl, triazolyl (e.g., 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl, etc.) tetrazolyl (e.g., 1H-tetrazolyl, 2H-tetrazolyl, etc.), etc.; unsaturated condensed heterocyclic group containing 1 to 5 nitrogen atoms, for example, indolyl, isoindolyl, indoliziny, benzimidazolyl, quinolyl, isoquinolyl,

indazolyl, benzotriazolyl, tetrazolopyridazinyl (e.g., tetrazolo [1,5-b]pyridazinyl, etc.), etc.; unsaturated 3 to 6-membered heteromonocyclic group containing an oxygen atom, for example, pyranyl, furyl, etc.; unsaturated 5 to 7-membered heteromonocyclic group containing a sulfur atom, for example, thienyl, etc;

5 unsaturated 5 to 7 membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, oxazolyl, isoxazolyl, oxadiazolyl (e.g., 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, etc.) etc.; unsaturated condensed heterocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms (e.g., benzoxazolyl, benzoxadiazolyl, etc.); unsaturated 5 to 7-membered

10 heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, for example, thiazolyl, thiadiazolyl (e.g., 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, etc.) etc.; unsaturated condensed heterocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms (e.g., benzothiazolyl, benzothiadiazolyl, etc.) and the like. The term also embraces radicals where heterocyclic radicals are

15 fused with aryl radicals. Examples of such fused bicyclic radicals include benzofuryl, benzothieryl, and the like. Said "heterocyclyl" radicals may also be substituted at a substitutable position with one or more substituents selected independently from alkyl, aralkyl, alkoxyalkyl, alkylaminoalkyl, carboxyalkyl, alkoxy carbonylalkyl, aminocarbonylalkyl, alkoxy, aralkoxy, alkylaminoalkoxy, aminocarboxy, alkylaminocarboxy, aralkylaminocarboxy, amino, halo, nitro,

20 alkylamino, alkylcarbonylamino, arylcarbonylamino, alkoxy carbonylamino, aralkoxy carbonylamino, aminocarbonylamino, alkylaminocarbonylamino, alkylsulfonylamino, arylsulfonylamino, acyl, cyano, carboxy, aminocarbonyl, alkoxy carbonyl and aralkoxy carbonyl. More preferred heteroaryl radicals include 5

25 to 6-membered heteroaryl radicals.

The term "cycloalkyl" embraces radicals having 3 to 10 carbon atoms. More preferred cycloalkyl radicals are "lower cycloalkyl" radicals having 3 to 7 carbon atoms. Examples include radicals such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

30 The term "sulfonyl", whether used alone or linked to other terms such as alkylsulfonyl, denotes respectively divalent radicals $\text{-SO}_2\text{-}$. "Alkylsulfonyl"

embraces alkyl radicals attached to a sulfonyl radical, where alkyl is defined as above. More preferred alkylsulfonyl radicals are "lower alkylsulfonyl" radicals having one to six carbon atoms. Examples of such lower alkylsulfonyl radicals include methylsulfonyl, ethylsulfonyl and propylsulfonyl. The "alkylsulfonyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide "haloalkylsulfonyl" radicals. More preferred haloalkylsulfonyl radicals are "lower haloalkylsulfonyl" radicals having one or more halo atoms attached to lower alkylsulfonyl radicals as described above. Examples of such lower haloalkylsulfonyl radicals include fluoromethylsulfonyl, trifluoromethylsulfonyl, and chloromethylsulfonyl. The term "arylsulfonyl" embraces aryl radicals as defined above, attached to a sulfonyl radical. Examples of such radicals include phenylsulfonyl. The terms "sulfamyl", "aminosulfonyl" and "sulfonamidyl" denotes $\text{NH}_2\text{O}_2\text{S}$.

The terms "carboxy" or "carboxyl", whether used alone or with other terms, such as "carboxyalkyl", denotes $-\text{CO}_2\text{H}$.

The term "carbonyl", whether used alone or with other terms, such as "alkoxycarbonyl", denotes $-(\text{C}=\text{O})-$. The term "alkoxycarbonyl" means a radical containing an alkoxy radical, as defined above, attached via an oxygen atom to a carbonyl radical. Preferably, "lower alkoxycarbonyl" embraces alkoxy radicals having one to six carbon atoms. Examples of such "lower alkoxycarbonyl" ester radicals include substituted and unsubstituted methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, and hexyloxycarbonyl.

The term "aralkyl" embraces aryl-substituted alkyl radicals. Preferable aralkyl radicals are "lower aralkyl" radicals having aryl radicals attached to alkyl radicals having 1 to 6 carbon atoms. Examples of such radicals include benzyl, diphenylmethyl, triphenylmethyl, phenylethyl and diphenylethyl. The aryl in the aralkyl may be additionally substituted as described above.

The term "aralkenyl" embraces aryl-substituted alkenyl radicals. Preferable aralkenyl radicals are "lower phenylalkenyl" radicals having phenyl radicals attached to alkenyl radicals having 1 to 6 carbon atoms. Examples of such radicals include phenylethenyl and phenylpropenyl. The aryl in said aralkyl may be additionally

substituted as described above. The terms benzyl and phenylmethyl are interchangeable.

The term "alkylcarbonyl" includes radicals having alkyl radicals as defined above, attached to a carbonyl radical. More preferred alkylcarbonyl radicals are "lower alkylcarbonyl" radicals having one to six carbon atoms. Examples of such radicals include methylcarbonyl and ethylcarbonyl.

The term "carboxyalkyl" embraces radicals having a carboxy radical as defined above, attached to an alkyl radical. The alkanoyl radicals may be substituted or unsubstituted, such as formyl, acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl, trifluoroacetyl or the like, in which the preferable one is formyl, acetyl, propionyl or trifluoroacetyl.

The term "heterocyclylalkyl" embraces heterocyclic-substituted alkyl radicals. More preferred heterocyclylalkyl radicals are "lower heterocyclylalkyl" radicals having 5 to 6 membered heterocyclyl radicals attached to lower alkyl radicals having 1 to 6 carbon atoms. Examples of such radicals include pyrrolidinylmethyl, piperidinylmethyl, morpholinylmethyl, piperazinylmethyl, oxazolylmethyl, oxazolylethyl, oxazolinylmethyl, oxazoliny lethyl, indoly lethyl, indoly lmethyl, pyridylmethyl, quinoly lmethyl, thienylmethyl, fury lethyl and quinoly lethyl. The heterocyclic in said heterocyclylalkyl may be additionally substituted as described above.

The term "aryloxy" embraces aryl radicals, as defined above, attached to an oxygen atom. The aryl in said aryloxy may be additionally substituted as described above. Examples of such radicals include phenoxy.

The term "aralkoxy" embraces oxy-containing aralkyl radicals attached through an oxygen atom to other radicals. The "aralkoxy" radical may be further substituted on the aryl ring portion of the radical.

The term "alkylamino" denotes amino groups which have been substituted with 1 or 2 alkyl radicals. More preferred alkylamino radicals are "lower alkylamino" having alkyl radicals of 1 to 6 carbon atoms attached to the nitrogen atom of an amine. Suitable "lower alkylamino" may be mono or dialkylamino such

as N-methylamino, N-ethylamino, N,N-dimethylamino, N,N-diethylamino or the like.

The term "alkylaminoalkyl" denotes alkylamino groups, as defined above, attached to an alkyl radical. More preferred alkylaminoalkyl radicals are "lower alkylaminoalkyl" having 1 to 6 carbon atoms attached to a lower aminoalkyl radical as described above. Suitable "lower alkylaminoalkyl" may be mono or dialkylaminoalkyl radicals such as N-methylaminomethyl, N-ethylaminomethyl, N,N-dimethylaminomethyl, N,N-dimethylaminoethyl, N,N-dimethylaminopropyl or the like.

The term "dialkylaminoalkyl" also includes radicals where the bridging alkyl moiety is optionally substituted with alkylsulfonyl, alkoxy, aralkoxy, heterocyclyl, and aryl.

The term "alkylaminoalkoxy" denotes alkylamino groups, as defined above, attached to an alkoxy radical. Suitable "alkylaminoalkoxy" may be mono or dialkylaminoalkoxy radicals such as N-methylaminomethoxy, N-ethylaminomethoxy, N,N-dimethylaminomethoxy, N,N-dimethylaminoethoxy, N,N-dimethylaminopropoxy or the like.

The term "alkylaminocarbonyl" embraces alkylamino radicals, as described above, to a carbonyl radical. More preferred alkylaminocarbonyl radicals are "lower alkylaminocarbonyl" having lower alkylamino radicals, as described above, attached to a carbonyl radical. Examples of such radicals include N-methylaminocarbonyl and N,N-dimethylaminocarbonyl.

The term "arylamino" denotes amino groups which have been substituted with 1 or 2 aryl radicals, such as N-phenylamino. The "arylamino" radicals may be further substituted on the aryl ring portion of the radical.

The terms "N-arylaminoalkyl" and "N-aryl-N-alkyl-aminoalkyl" denote amino groups which have been substituted with one aryl radical or one aryl and one alkyl radical, respectively, and having the amino group attached to an alkyl radical. More preferred arylaminoalkyl radicals are "lower arylaminoalkyl" having the arylamino radical attached to 1 to 6 carbon atoms. Examples of such radicals include N-phenylaminomethyl and N-phenyl-N-methylaminomethyl.

The term "aminocarbonyl" denotes an amide group of the formula -
C(=O)NH₂. The term "aminocarbonylalkyl" denotes an aminocarbonyl group
attached to an alkyl radical. More preferred are "lower aminocarbonylalkyl" having
lower aminocarbonyl radicals as described above attached to alkyl of one to six
carbon atoms. The term "alkylaminocarbonylalkyl" denotes an aminocarbonyl
group which has been substituted with one or two alkyl radicals and attached to an
alkyl radical. More preferred are "lower alkylaminocarbonylalkyl" having lower
alkylaminocarbonyl radicals as described above attached to alkyl radicals of one to
six carbon atoms.

The term "aryloxy" embraces aryl radicals attached to a divalent oxygen
atom, that is, to form monoaryloxy and diaryloxy radicals. The more preferred
aryloxy radicals are "lower aryloxy". An example includes phenoxy.

"Amino acid residue" means any of the naturally occurring alpha-, beta- and
gamma-amino carboxylic acids, including their D and L optical isomers and racemic
mixtures thereof, synthetic amino acids, and derivatives of these natural and
synthetic amino acids. The amino acid residue is bonded through a nitrogen of the
amino acid. The naturally occurring amino acids which can be incorporated in the
present invention include, but are not limited to, alanine, arginine, asparagine,
aspartic acid, cysteine, cystine, glutamic acid, glutamine, glycine, histidine,
isoleucine, leucine, lysine, methionine, ornithine, phenylalanine, proline, serine,
threonine, thyroxin, tryptophan, tyrosine, valine, β -alanine, and γ -aminobutyric
acid. Derivatives of amino acids which can be incorporated in the present invention
include, but are not limited to, amino acids having protected and modified
carboxylic acids, including acid esters and amides, protected amines, and substituted
phenyl rings, including but not limited to alkyl, alkoxy and halo substituted tyrosine
and phenylalanine.

The present invention comprises a pharmaceutical composition comprising a
therapeutically-effective amount of a compound of Formula II in association with at
least one pharmaceutically-acceptable carrier, adjuvant or diluent.

The present invention also comprises a method of therapeutic and
prophylactic treatment of viral infections, particularly herpetoviridae infection, in a

subject, the method comprising treating the subject having such herpes infection a therapeutically-effective amount of a compound of Formula II.

The present invention also comprises a method of inhibiting a viral protease, the method comprising administering a therapeutically-effective amount of a compound of Formula II.

Also included in the family of compounds of Formula II are the stereoisomers and tautomers thereof. Compounds of the present invention can possess one or more asymmetric carbon atoms and are thus capable of existing in the form of optical isomers as well as in the form of racemic or non-racemic mixtures thereof. Accordingly, some of the compounds of this invention may be present in racemic mixtures which are also included in this invention. The optical isomers can be obtained by resolution of the racemic mixtures according to conventional processes, for example by formation of diastereoisomeric salts by treatment with an optically active acid or base. Examples of appropriate acids are tartaric, diacetyltartaric, dibenzoyltartaric, ditoluoyltartaric and camphorsulfonic acid and then separation of the mixture of diastereoisomers by crystallization followed by liberation of the optically active bases from these salts.

A different process for separation of optical isomers involves the use of a chiral chromatography column optimally chosen to maximize the separation of the enantiomers. Still another available method involves synthesis of covalent diastereoisomeric molecules by reacting an amine functionality of precursors to compounds of Formula II with an optically pure acid in an activated form or an optically pure acid in an activated form or an optically pure isocyanate.

Alternatively, diastereomeric derivatives can be prepared by reacting a carboxyl functionality of precursors to compounds of Formula II with an optically pure amine base. The synthesized diastereoisomers can be separated by conventional means such as chromatography, distillation, crystallization or sublimation, and then hydrolyzed to deliver the enantiomerically pure compound. The optically active compounds of Formula II can likewise be obtained by utilizing optically active starting materials. These isomers may be in the form of a free acid, a free base, an ester or a salt. Additional methods for resolving optical isomers, known to those

skilled in the art may be used, for example, those discussed by Jaques et al. in *Enantiomers, Racemates, and Resolutions*, John Wiley and Sons, New York (1981).

Also included in the family of compounds of Formula II are the pharmaceutically-acceptable salts thereof. The term "pharmaceutically-acceptable salts" embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts of compounds of Formula II may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, examples of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, salicylic, *p*-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethylsulfonic, benzenesulfonic, pantothenic, toluenesulfonic, 2-hydroxyethanesulfonic, sulfanilic, stearic, cyclohexylaminosulfonic, algenic, β -hydroxybutyric, galactaric and galacturonic acid. Suitable pharmaceutically-acceptable base addition salts of compounds of Formula II include metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from N,N'-dibenzylethylenediamine, choline, chlorprocaine, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. All of these salts may be prepared by conventional means from the corresponding compound of Formula II by reacting, for example, the appropriate acid or base with the compound of Formula II.

Also embraced within this invention is a class of pharmaceutical compositions comprising one or more compounds of Formula II in association with one or more non-toxic, pharmaceutically-acceptable carriers and/or diluents and/or adjuvants (collectively referred to herein as "carrier" materials) and, if desired, other active

ingredients. The compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. The compounds and composition may, for example, be administered orally, intravascularly, intraperitoneally, subcutaneously, intramuscularly or topically.

For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are tablets or capsules. The active ingredient may also be administered by injection as a composition wherein, for example, saline, dextrose or water may be used as a suitable carrier.

The amount of therapeutically active compound that is administered and the dosage regimen for treating a disease condition with the compounds and/or compositions of this invention depends on a variety of factors, including the age, weight, sex and medical condition of the subject, the severity of the disease, the route and frequency of administration, and the particular compound employed, and thus may vary widely. The pharmaceutical compositions may contain active ingredient in the range of about 0.1 to 2000 mg, preferably in the range of about 0.5 to 500 mg and most preferably between about 1 and 100 mg. A daily dose of about 0.01 to 100 mg/kg body weight, preferably between about 0.1 and about 50 mg/kg body weight and most preferably between about 1 to 20 mg/kg body weight, may be appropriate. The daily dose can be administered in one to four doses per day.

For therapeutic purposes, the compounds of this invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered *per os*, the compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanoic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may contain a controlled-

release formulation as may be provided in a dispersion of active compound in hydroxypropylmethyl cellulose. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut, oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.

For infections of the eye or other external tissues, e.g., mouth and skin, the formulations are preferably applied as a topical ointment or cream containing the active ingredient(s) in a total amount of, for example, 0.075 to 30 % w/w, preferably 0.2 to 20 % w/w and most preferably 0.4 to 15 % w/w. When formulated in an ointment, the active ingredients may be employed with either paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base. If desired, the aqueous phase of the cream base may include, for example, at least 30 % w/w of a polyhydric alcohol such as propylene glycol, butane-1,3-diol, mannitol, sorbitol, glycerol, polyethylene glycol and mixtures thereof. The topical formulation may desirably include a compound which enhances absorption or penetration of the active ingredient through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethylsulfoxide and related analogs. The compounds of this invention can also be administered by a transdermal device. Preferably topical administration will be accomplished using a patch either of the reservoir and porous membrane type or of a solid matrix variety. In either case, the active agent is delivered continuously from the reservoir or microcapsules through a membrane into the active agent permeable adhesive, which is in contact with the skin or mucosa of the recipient. If the active agent is absorbed through the skin, a controlled and predetermined flow of the active agent is administered to the

recipient. In the case of microcapsules, the encapsulating agent may also function as the membrane.

The oily phase of the emulsions of this invention may be constituted from known ingredients in a known manner. While the phase may comprise merely an emulsifier, it may comprise a mixture of at least one emulsifier with a fat or an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make-up the so-called emulsifying wax, and the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations. Emulsifiers and emulsion stabilizers suitable for use in the formulation of the present invention include Tween 60, Span 80, cetostearyl alcohol, myristyl alcohol, glyceryl monostearate, and sodium lauryl sulfate, among others.

The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties, since the solubility of the active compound in most oils likely to be used in pharmaceutical emulsion formulations is very low. Thus, the cream should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic alkyl esters such as di-isoadipate, isocetyl stearate, propylene glycol diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-thylhexyl palmitate or a blend of branched chain esters may be used. These may be used alone or in combination depending on the properties required. Alternatively, high melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils can be used.

Formulations suitable for topical administration to the eye also include eye drops wherein the active ingredients are dissolved or suspended in suitable carrier, especially an aqueous solvent for the active ingredients. The antiviral active ingredients are preferably present in such formulations in a concentration of 0.5 to 20 %, advantageously 0.5 to 10 % and particularly about 1.5 % w/w.





Examples





5 The following examples contain detailed descriptions of the methods of preparation of compounds of Formula II. These detailed descriptions fall within the scope, and serve to exemplify, the above described General Synthetic Procedures which form part of the invention. These detailed descriptions are presented for illustrative purposes only and are not intended as a restriction on the scope of the invention. All parts are by weight and temperatures are in Degrees centigrade unless otherwise indicated.





The following abbreviations are used:





EtOAc	ethyl acetate
HCl	hydrochloric acid
DMSO	dimethylsulfoxide
d ₆ -DMSO	deuterated dimethylsulfoxide
CDCl ₃	deuterated chloroform
CHCl ₃	chloroform
CD ₃ OD	deuterated methanol
Et ₂ O	diethyl ether
MgSO ₄	magnesium sulfate
H ₂ SO ₄	sulfuric acid
NaHCO ₃	sodium bicarbonate
KHSO ₄	potassium hydrogen sulfate
NMM	N-methylmorpholine
DMF	dimethylformamide
DMAP	4-dimethylaminopyridine
CDI	carbonyldiimidazole
NaOH	sodium hydroxide
KOH	potassium hydroxide
LiOH	lithium hydroxide
Pd(OH) ₂ /C	palladium hydroxide on carbon
Pd/C	palladium on carbon
EDC	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide-HCl
BOC	<i>tert</i> -butyloxycarbonyl
TLC	thin layer chromatography
MeOH	methanol
KI	potassium iodide
CH ₂ Cl ₂	methylene chloride



The following is a list of inventive compounds and comparative compounds.





Compound	Formula
1	
2	
3	
4	



Compound	Formula
5	 <chem>CC1=C(C2=CC=CC=C2N)CCN(C)C3=NC4=C(C(=O)O3)C=C(C=C4)NC(=O)OC(C)(C)C</chem>
6	 <chem>CC1=C(C2=CC=CC=C2N3CCOCC3)C=NC4=C(C(=O)O4)C=C(C=C4)NC(=O)OC(C)(C)C</chem>
7	 <chem>CC1=C(C2=CC=CC=C2N(C)CC=C)C=NC3=C(C(=O)O3)C=C(C=C4)NC(=O)OC(C)(C)C</chem>
8	 <chem>CC1=C(C2=CC=CC=C2N3CCCCC3)C=NC4=C(C(=O)O4)C=C(C=C4)NC(=O)OC(C)(C)C</chem>

Compound	Formula
9	 <chem>CC(C)N1C(=O)OC2=CC=C(C=C2N1)NC(=O)OCCc3ccccc3</chem>
10	 <chem>CC(C)N1C(=O)OC2=CC=C(C=C2N1)NC(=O)OC(C)(C)C</chem>
11	 <chem>C1CCN2C(=O)OC3=CC=C(C=C3N2C1)NC(=O)OC(C)(C)C</chem>
12	 <chem>CC(C)N1C(=O)OC2=CC=C(C=C2N1)NC(=O)C3=CC(=CC(=C3)F)F</chem>

Compound	Formula
13	 <chem>CC1=CC=C(C(=C1)C2=NC(=C3C(=O)N(C2)C(=O)OC(C)(C)C)C3)N(C4CC4)CC#N</chem>
14	 <chem>CC1=CC=C(C(=C1)C2=NC(=C3C(=O)N(C2)C(=O)OC(C)(C)C)C3)N(C4CCN(C4)C(=O)c5ccoc5)</chem>
15	 <chem>CC1=CC=C(C(=C1)C2=NC(=C3C(=O)N(C2)C(=O)OC(C)(C)C)C3)N(C4CCN(C4)S(=O)(=O)c5ccccc5)</chem>
16	 <chem>CC1=CC=C(C(=C1)C2=NC(=C3C(=O)N(C2)C(=O)OC(C)(C)C)C3)N(C4CCN(C4)C(=O)C)</chem>

Compound	Formula
17	
18	

Comparative Compound	Formula
CC 1	
CC 2	
CC 3	
CC 4	

Comparative Compound	Formula
CC 5	
CC 6	

Example 1**Compound 1**

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[methyl(phenylmethyl)amino]-4H-3,1-benzoxazin-4-one

A. Preparation of 3-[[[(1,1-dimethylethoxy)carbonyl]amino]-2-methyl-6-[[[methyl(phenylmethyl)-amino]carbonyl]amino]benzoic acid, 2-(trimethylsilyl)ethyl ester.

To a solution of 6-amino-3-[[[(1,1-dimethylethoxy)carbonyl]amino]-2-methylbenzoic acid, 2-(trimethylsilyl)ethyl ester (0.11 mmol in 3ml of CH_2Cl_2), was added a solution of p-Nitrophenyl chloroformate (0.1 mmol, 20.2 mg) in CH_2Cl_2 (0.2 ml). After stirring at room temperature for 3 hours, resulting solution was washed with 1N-HCl, water, and dried over MgSO_4 . To this solution of activated carbamate, methyl(phenylmethyl)amine (0.016 ml, 0.12 mmol) was added. After stirring at room temperature for 15 hours, a solution of tetrafluorophthalic anhydride (15 mg, 0.07 mmol) in CH_2Cl_2 (0.7 ml) was added and stirred for 3 hours, followed by polyamine resin prepared by the method of L. Flynn et al. (*J. Amer. Chem. Soc.*, 119, 4874-4881 (1997)) (200 mg, 0.6 mmol). After 1 hour, the

5 mixture was filtrated and concentrated, dissolved in CH_2Cl_2 again, Amberlyst A-21 (50 mg) was added, stirred for 1 hour. Resin was removed with filtration, the solution was concentrated to afford white crystal (38.7 mg). $^1\text{H-NMR}$ (270MHz, CDCl_3) δ 0.06(s, 9H), 1.0-1.1(m, 2H), 1.50(s, 9H), 2.28(s, 3H), 2.99(s, 3H), 4.3-4.4(m, 2H), 4.59(s, 2H), 6.15(br.s, 1H), 7.2-7.3(m, 5H), 7.67(br.d, 1H, $J=9\text{Hz}$), 8.01(d, 1H, $J=9\text{Hz}$), 8.25(s, 1H), HPLC(A) retention time=26.8min.

B. Preparation of 3-[[[(1,1-dimethylethoxy)carbonyl]amino]-2-methyl-6-[[[methyl(phenylmethyl)-amino]carbonyl]amino]benzoic acid.

10 To a solution of the product of step A(38.7 mg) in 3 ml of THF was added a solution of TBAF in THF (1.0 M, 0.12 ml, 0.12 mmol) and stirred at room temperature for 1 hour. Amberlyst A-15 H form (200 mg), Amberlyst A-15 calcium form (200 mg) were added, shaken overnight, filtrated, and concentrated to afford white crystal (26.0 mg). $^1\text{H-NMR}$ (270MHz, CD_3OD) δ 1.42(s, 9H), 2.24(s, 3H), 2.89(s, 3H), 4.50(s, 2H), 7.1-7.3(m, 6H), 7.64(d, 1H, $J=9\text{Hz}$), HPLC(A) retention time=15.4 min.

15 C. 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[[methyl(phenylmethyl)-amino]-4H-3,1-benzoxazin-4-one

20 To a solution of the product of step B(26.0 mg) in 4 ml of DMF was added P-EDC (300 mg, 0.3 mmol), stirred for 2 hours. Filtrated, concentrated to afford white crystal (21.5mg). $^1\text{H-NMR}$ (270MHz, CDCl_3) δ 1.51(s, 9H), 2.65(s, 3H), 3.08(s, 3H), 4.74(s, 2H), 6.26(br.s, 1H), 7.12(d, 1H, $J=9\text{Hz}$), 7.2-7.4(m, 5H), 7.83(br.d, 1H), HPLC(A) retention time=24.7min, $\text{MS}(\text{MH}^+)=396$.

A similar method was used to prepare the following compounds

Compound 3

25 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[[methyl[(4-methoxyphenyl)methyl]amino]-4H-3,1-benzoxazin-4-one. $^1\text{H-NMR}$ (270MHz, CDCl_3) δ 1.51(s, 9H), 2.65(s, 3H), 3.06(s, 3H), 3.80(s, 3H), 4.67(s, 2H), 6.23(br.s,

1H), 6.87(d, 2H, J=9Hz), 7.12(d, 1H, J=9Hz), 7.25(d, 2H, J=9Hz), 7.83(br.d, 1H),
HPLC(B) retention time=8.26 min, MS(MH+)=426.

Compound 8

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[1-(1,2,3,6-tetrahydro-
pyridyl)]-4H-3,1-benzoxazin-4-one.

HPLC(A) retention time=18.4 min, MS(MH+)=358.

Compound 10

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[isopropyl(methyl)amino]-
4H-3,1-benzoxazin-4-one.

HPLC(A) retention time=22.9 min.

Compound 14

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[(4-furoyl)piperazinyl]-4H-
3,1-benzoxazin-4-one.

HPLC(A) retention time=20.0 min.

Compound 13

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[(2-cyanoethyl)-
(cyclopropyl)amino]-4H-3,1-benzoxazin-4-one.

HPLC(A) retention time=21.0 min.

Compound 17

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[(4-thenoyl)piperazinyl]-4H-
3,1-benzoxazin-4-one.

HPLC(B) retention time=7.02 min.

Compound 15

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[(4-benzenesulfonyl)-
piperazinyl]-4H-3,1-benzoxazin-4-one.

HPLC(B) retention time=7.88 min.

Example 2

Compound 2

6-(benzoylamino)-5-methyl-2-[methyl(phenylmethyl)amino]-
4H-3,1-benzoxazin-4-one

A. Preparation of 3-amino-2-methyl-6-[[[methyl(phenylmethyl)amino]-carbonyl]amino]-benzoic acid, 2-(trimethylsilyl)ethyl ester.

To a solution of the product of example 1-step A (40.2 mg) in 1,4-dioxane (2 ml) was added 4N-HCl solution in 1,4-dioxane (2 ml). After stirred at room temperature for 2 hours, the reaction mixture was concentrated. CH_2Cl_2 (4 ml) and DMF (0.5 ml) was added, dissolved, and Amberlyst A-21 (100 mg) was added. Shaken for 1 hour, the resin was filtered off, concentrated to afford 35.7 mg of product. $^1\text{H-NMR}$ (270MHz, CDCl_3) δ 0.06(s, 9H), 0.9-1.0(m, 2H), 2.18(s, 3H), 3.97(s, 3H), 4.2-4.3(m, 2H), 4.58(s, 2H), 6.79(d, 1H, $J=8.9\text{Hz}$), 7.2-7.4(m, 5H), 7.77(d, 1H, $J=8.9\text{Hz}$), 8.14(s, 1H), HPLC(A) retention time=23.4min.

B. Preparation of 3-(benzoylamino)-2-methyl-6-[[[methyl(phenylmethyl)amino]-carbonyl]amino]-benzoic acid, 2-(trimethylsilyl)ethyl ester.

To a solution of the product of step A (35.7 mg) in 2 ml of CH_2Cl_2 , were added pyridine (0.011 ml, 0.14 mmol), benzoyl chloride (0.014 ml, 0.12 mmol). After stirring at room temperature for 14 hours, polyamine resin (200 mg, 0.6 mmol) was added, stirred for 2 hours. Filtration and concentration afforded 32.3 mg of the product. $^1\text{H-NMR}$ (270MHz, CDCl_3) δ 0.06(s, 9H), 1.0-1.1(m, 2H), 2.33(s, 3H), 2.98(s, 3H), 4.2-4.3(m, 2H), 4.57(s, 2H), 7.2-7.6(m, 9H), 7.8-7.9(m, 3H), 8.07(d, 1H, $J=9\text{Hz}$), 8.84(s, 1H), HPLC(A) retention time=25.4min.

C. Preparation of 3-(benzoylamino)-2-methyl-6-[[[methyl(phenylmethyl)amino]-carbonyl]amino]-benzoic acid.

To a solution of the product of step B (32.3 mg) in 3ml of THF was added a solution of TBAF in THF (1.0 M, 0.12 ml, 0.12 mmol) and stirred at room temperature for 1 hour. Amberlyst A-15 H form (200 mg), Amberlyst A-15 calcium form (200 mg) were added, shaken overnight, filtrated, and concentrated to afford white crystal (25.4 mg). ¹H-NMR (270MHz, CDCl₃) δ 2.19(s, 3H), 2.90(s, 3H), 4.52(s, 2H), 7.2-7.5(m, 7H), 7.77(d, 1H, J=9Hz), 7.9(m, 2H), 8.58(s, 1H), 8.96(br.s, 1H).

D. 6-(benzoylamino)-5-methyl-2-[methyl(phenylmethyl)amino]-4H-3,1-benzoxazin-4-one.

To a solution of the product of step C (25 mg) in 4ml of DMF was added P-EDC (300 mg, 0.3 mmol), stirred for 2 hours. Filtrated, concentrated to afford white crystal (15.8 mg). ¹H-NMR (270MHz, CDCl₃) δ 2.69(s, 3h), 3.01(s, 3H), 4.75(s, 2H), 7.16(d, 1H, J=9Hz), 7.2-7.6(m, 8H), 7.8-8.0(m, 4H), HPLC(A) retention time=22.4 min., MS(MH⁺)=400.

Compound 9

6-[(Phenylmethoxyacetyl)amino]-5-methyl-2-(diisopropylamino)-4H-3,1-benzoxazin-4-one.

HPLC(A) retention time=24.1 min.

Compound 12

6-[(2,4,6-Trifluorobenzoyl)amino]-5-methyl-2(diisopropylamino)-4H-3,1-benzoxazin-4-one.

HPLC(A) retention time=23.3 min.

Example 3**Compound 4**

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[[[4-(dimethylamino)phenyl]methyl]methylamino]-4H-3,1-benzoxazin-4-one.

- 5 A. Preparation of 3-[[[(1,1-dimethylethoxy)carbonyl]amino]-2-methyl-6-[[[N-[[4-(dimethylamino)phenyl]methyl]methylamino]carbonyl]amino]benzoic acid, 2-(trimethylsilyl)ethyl ester.

Proceeding in a manner similar to example 1A, substituting methyl(phenylmethyl)amine with N-[[4-(dimethylamino)phenyl]methyl]methylamine. ¹H-NMR(270 MHz, CDCl₃) δ 0.06(s, 9H), 1.0-1.1(m, 2H), 1.50(s, 9H), 2.28(s, 3H), 2.93(s, 9H), 4.2-4.3(m, 2H), 4.48(s, 2H), 6.11(br.s, 1H), 6.70(d, 2H, J=9Hz), 7.19(d, 2H, J=9Hz), 7.58(br.d, 1H), 8.06(d, 1H, J=9Hz), 8.62(br. S, 1H).

- 10 B. Preparation of 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[[[4-(dimethylamino)phenyl]methyl]methylamino]-4H-3,1-benzoxazin-4-one.

15 To a solution of the product of step A (521 mg, 0.935 mmol) in 10 ml of THF, was added a solution of tetrabutylammonium fluoride in THF (1.0 M, 1.12 ml, 1.12 mmol). Stirred at room temperature for 45 min., acetyl chloride (0.12 ml, 1.7 mmol) was added, and stirred for 1.5 hours. Concentrated, purified over silica gel, washed with saturated NaHCO₃ aqueous solution to afford white amorphous solid (241 mg). ¹H-NMR (270MHz, CDCl₃) δ 1.51(s, 9H), 2.65(s, 3H), 2.93(s, 6H), 3.11(s, 3H), 4.63(s, 2H), 6.21(br.s, 1H), 6.69(d, 2H, J=9Hz), 7.12(d, 1H, J=9Hz), 7.19 (d, 2H, J=9Hz), 7.82(br.d, 1H, J=9Hz), HPLC(B) retention time=8.62 min, MS(MH⁺)=439.

Compound 18

20 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[methyl(2-pyridylmethyl)-amino]-4H-3,1-benzoxazin-4-one.
HPLC(B) retention time=6.71 min., MS(MH⁺)=397.

Compound 5

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[methyl[2-(3-indolyl)ethyl]amino]-4H-3,1-benzoxazin-4-one.

¹H-NMR (270MHz, CDCl₃) δ 1.52(s, 9H), 2.65(s, 3H), 3.1(m, 5H), 3.82(t, 2H, J=7Hz), 6.21(br.s, 1H), 7.0-7.4(m, 4H), 7.7-8.1(m, 3H). HPLC(B) retention time=8.08min., MS(MH⁺)=449.

Compound 6

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-(4-morpholyl)-4H-3,1-benzoxazin-4-one.

¹H-NMR (270MHz, CDCl₃) δ 1.52(s, 9H), 2.65(s, 3H), 3.72(d, 4H, J=3Hz), 3.75(d, 4H, J=3Hz), 6.25(br.s, 1H), 7.19(d, 1H, J=9Hz), 7.86(br.d, 1H, J=9Hz). HPLC(B) retention time=5.84min., MS(MH⁺)=362.

Compound 7

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[allyl(methyl)amino]-4H-3,1-benzoxazin-4-one.

¹H-NMR (270MHz, CDCl₃) δ 1.41(s, 9H), 2.44(s, 3H), 2.99(s, 3H), 4.06(d, 2H, J=5Hz), 5.1-5.3(m, 2H), 5.7-6.0(m, 1H), 6.98(d, 1H, J=9Hz), 7.44(d, 1H, J=9Hz), 8.68(br.s, 1H). HPLC(A) retention time=21.88min., MS(MH⁺)=346.

Compound 11

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-(1-decahydroquinolyl)-4H-3,1-benzoxazin-4-one.

HPLC(B) retention time=8.63min., MS(MH⁺)=414.

Compound 16

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[4-(1-acetylpiperadiny)]-4H-3,1-benzoxazin-4-one.

HPLC(B) retention time=4.98min., MS(MH⁺)=403.

Biological Evaluation

The compounds of this invention exhibited antiviral activity as indicated by inhibition *in vitro* of herpesvirus protease and HCMV infectivity. The antiviral activity of the compounds of this invention illustrated in the Examples was determined by the following methods.

Enzymatic assay for HSV-1 protease (assemblin) inhibition

FP Assay

Assemblin protease activity was determined using fluorescence polarization (FP). The fluorescent substrate was biotin-gamma-aminobutyrate-HTYLQASERFRIK-DTAF, based on the HSV-1 release cleavage site. Incubation of this substrate with assemblin resulted in cleavage between alanine and serine. A change in molecular size of fluorescent substrate molecule, which was increased using avidin as a reaction stop reagent, allowed cleavage to be measured by FP. Potential protease inhibitors were dissolved in DMF and then diluted 5-fold in assay buffer. 6.5uL were added to the wells of a 96-well plate (U-bottom 96-well black plate, Dynatech or Costar) which was previously blocked using blocking solution (10mM Tris-HCl pH 8.0, 150mM NaCl, 0.05% Tween 20, 1mg/mL BSA). The enzyme was diluted to 21.3 ug/mL in assay buffer (1M NaCitrate, 50mM NaPhosphate, pH 7.4, 100mM NaCl, 20% glycerol, 2mM TECP) and 48.5uL were added to each well. Following a 30 minute incubation at room temperature, 10uL of 62.5ug/mL substrate were added. After about 90 minute incubation at room temperature, 50uL of 2mg/mL avidin in phosphate buffer (50mM Naphosphate, 100mM NaCl, pH 7.4) were added to each well. The plates were read by a fluorescence polarization plate reader. Inhibitor potency was determined by comparison with incubations lacking inhibitor.

HPLC assay

Assays were performed with the peptide substrate H-His-Thr-Tyr-Leu-Gln-Ala-Ser-Glu-Lys-Phe-Lys-Met-Trp-Gly-NH₂ (Bachem). This substrate is a HSV-1-amide UL26 Open Reading Frame (242-255) and is derived from the release site of HSV-1 protease. HSV-1 protease cleaves between alanine and serine. The product

SEKFKMWG was quantified on HPLC using fluorescence detection of tryptophan residue. Enzyme was diluted to 4.3ug/mL in assay buffer (1M NaCitrate, 50mM NaPhosphate, pH 7.3, 100mM NaCl, 20% glycerol, 2mM TCEP) and 48.5uL were added to the tubes. Potential protease inhibitors were dissolved in DMF and then diluted 10-fold in assay buffer. 6.5uL of inhibitor solution were added to each tube. Following a 30-minute incubation at room temperature, 10uL of substrate in phosphate buffer (50mM Naphosphate, 100mM NaCl, pH 7.4) were added to all tubes. The final concentration of the substrate in the reaction mixtures was 10uM. After a 15-minute incubation at room temperature, assays were quenched using 50uL of 50% TCA. Inhibitor potency was determined by comparison with incubations lacking inhibitor, which under these conditions gave about 20% cleavage of substrate.

Assay Components

Recombinant HSV-1 protease

HSV-1 protease was purified from baculovirus expressing a DNA construction encoding residues 1-288 of HSV-1 UL26 open reading frame and 32 heterologous amino acid. The construction also encoded six additional histidine residues at the amino terminus of the protease. These additional histidine residues provided an affinity ligand by which the protein was purified using Ni-NTA agarose gel (Qiagen). The purified protease was stored in stock solution (20mM HEPES buffer, pH 8.5, containing 20% (v/v) glycerol). This stock was diluted with assay buffer to adequate concentration of enzymatic assay.

Substrate

FP Assay

A specific substrate was synthesized based on the cleavage specificity of HSV-1 protease at the "release site" of the assembly protein (DiIanni, C.L., et al., *J. Biol. Chem.* 268, 2048, (1993)). The assembly protein release site has the sequence, HTYLQA*SEKFKMWG. The substrate used was biotin-gamma-aminobutyrate-HTYLQA*SERFRIK-DTAF which was prepared by standard peptide synthetic methods such as that described in Bodansky and Bodansky, "The

Practice of Peptide Synthesis" (1984), and was stored as a stock solution at 2.5mg/mL in DMF. This was diluted to 62.5ug/mL with phosphate buffer (50mM Naphosphate, 100mM NaCl, pH 7.4) just before use.

HPLC assay

5 The substrate was HSV-1-amide UL 26 Open Reading Frame (242-255), obtained from Bachem (Product No. M-2160).

Assay Buffer

10 An assay buffer (1M NaCitrate, 50mM NaPhosphate, pH 7.4, 100mM NaCl, 20% glycerol, 2mM TCEP) was used to dilute stock solutions of enzyme and inhibitors.

Antiviral and Cytotoxic Assay

These complimentary assays tested the ability of a compound to inhibit the production of new virus and the toxicity of the compound to the host cells. It was important that both assays be performed simultaneously in order to compare the results directly since toxicity may indirectly reduce viral replication.

Abbreviations:

DMEM - Dulbecco's Modified Eagle Medium; commercially available.

FBS - fetal bovine serum; commercially available and unknown factors necessary for growth of cells in culture.

20 HSV - herpes simplex virus.

Antiviral assay

25 The antiviral assay was estimated by plaque reduction assay performed by following methods. 1×10^5 of vero cells (African green monkey kidney cell) in 48-well plates were over-night cultured and the medium of this culture was replaced with 200 μ l of 2% FBS DMEM containing 2 \times the desired final concentration of test compounds or no compound as control. The cultures were added 50 μ l of 2% FBS DMEM containing with about 50 plaque forming units of HSV-1 and next 250 μ l of 2% FBS DMEM containing 1% methylcellulose. These infected cultures were incubated at 37°C, 5% CO₂ for 3 days until plaques was visible. The cells

were fixed and stained simultaneously with 0.025% crystalviolet in 5% formalin solution and plaque were counted. The concentration of test compound which conferred 50% inhibition of plaque formation compared to no compound control was interpolated from the observed data and defined as IC_{50} . Results are included in Table 2.

Cytotoxic assay

4×10^4 of vero cells in 96-wells were over-night cultured in 100 μ l of 10% FBS DMEM. These cultured cells were added 100 μ l of 10% FBS DMEM containing 50 μ M of test compounds or no compound as control. Cells were cultured incubated at 37°C, 5% CO₂ for 3 days. For measuring proliferation of the cells, cells were added 20 μ l of alamarBlue™ and incubated for 8 hours until the color of control was changed. And then the cells were read spectrophotometrically (absorbance at 570nm and 600nm) with BIO-RAD model 3550 microplate reader. The result was indicted as the ratio at 50 μ M compound concentration and no compound.

Chymotripsin Assay

The chymotripsin assay was modified from the method of Delmar, et al. (Anal. Biochem. 99, 316-320 (1979)). Bovine pancreas α -chymotripsin(type II, Sigma) was dissolved in 0.001 N HCl at 1 mg/ml and further diluted 1/1000 in assay buffer (0.1 M Tris, pH 7.8 containing 0.1 M CaCl₂) before use. 0.75 μ l of test compound in DMF (or DMF alone), 50 μ l of assay buffer and 50 μ l of enzyme were added to 96 wells plates, mixed and pre-incubated for 30 minutes at ambient temperature. Reaction was initiated by addition of 50 μ l of 0.2 mM N-succinyl-Ala-Ala-Pro-Phe-p-nitroanilide (Sigma; 2 mM in DMSO diluted 1/10 in assay buffer before use). The increase in absorbance at 405 nm was monitored for 3 minutes with BIO-RAD model 3550 microplate reader.

Human Leukocyte Elastase Assay

Human leukocyte elastase (HLE) (CALBIOCHEM) was dissolved in saline at 1 mg/ml and further diluted 1/10 in assay buffer (0.2 M Tris, pH 8.0) before use.

0.75 μ l of test compound in DMF (or DMF alone), 50 μ l of assay buffer and 50 μ l of enzyme were added to 96 well plate, mixed and pre-incubated for 30 minutes at ambient temperature. Reaction was initiated by addition of 50 μ l of 2.5 mM methoxysuccinyl-Ala-Ala-Pro-Val-p-nitroanilide (Sigma; 25 mM in DMSO diluted 1/10 in assay buffer before use). The increase in absorbance at 405 nm was monitored for 3 min with BIO-RAD model 3550 microplate reader.

HPLC

condition A

Column: YMC-Pack ODS A-312 6.0 \times 150mm

Eluent system:

Time(min)	0	20	30
MeCN(%)	5	100	100
10mMNaH ₂ PO ₄ aq.	95	0	0

Detection UV 254nm

Flow rate 1.0ml/min

condition B

Column: COSMOSIL 3C18 4.6 \times 50mm

Eluent system:

Time(min)	0	5	15
MeCN(%)	30	90	90
10mMNaH ₂ PO ₄ aq.	70	10	10

Detection UV 254nm


Flow rate 1.0ml/min.

COMPOUND	ENZYME IC50 (μ M)	ANTIVIRUS EC50 (μ M)	INHIBITION AT 50 μ M	
			HLE (%)	CHYMOTRYPSIN (%)
1	0.6	1.1	32	13
2	0.7	4.6	26	7
3	0.6	0.16	0	0
4	1.3	0.05	0	0
5	3.5	3.2	0	13.1
6	3.7	2.9	0	0
7	3.4	4.1	0	6.9
8	5	3.3	0	0
9	1.5	11.2	18.2	28.4
10	7.8	3.3	0	0
11	5.5	2.8	0	0
12	6.3	6.8	13.4	22.4
13	3.9	7.6	0	2.2
14	2	1.6	46.3	0
15	1.7	2.8	0	0
16	2.3	3.6	1.4	0
17	4.2	4.1	0	0
18	1.5	15.1	28.7	0
CC1	20.0	no data	no data	no data
CC2	3.2	7.5	101	101
CC3	3.2	48.9	100	100
CC4	no data	15.0	98	102
CC5	28.0	>50	100	98
CC6	11.8	>50	99	97

It will be apparent to those skilled in the art that various modifications and variations can be made in the compositions and methods of the present invention without departing from the spirit or scope of the invention. Thus, it is intended that the present invention cover the modifications and variations of this invention
5 provided they come within the scope of the appended claims and their equivalents.

WHAT IS CLAIMED IS:




1. A compound of Formula II:



II

wherein R^{28} is selected from amino optionally substituted with two radicals selected from alkyl, aralkyl, heterocyclalkyl, heterocycl, and aryl;


wherein R^{29} is selected from



wherein R^{30} is selected from alkyl, alkoxy, alkylamino, carboxyalkyl, alkoxyalkyl, alkylaminoalkyl, cycloalkyl, heterocycl, heterocyclalkyl, aralkyl, aralkoxy, aryloxy, cycloalkyloxy, arylamino, aralkenyl, heterocyclalkoxy, alkylaminoalkoxy, alkylaminoalkylamino, heterocyclalkylamino, N-aryl-N-alkylamino, and N-aralkylamino; wherein R^{31} is alkyl; wherein R^{32} is selected from alkyl and aryl; and wherein R^{33} is selected from hydrido, halo and alkyl;

or a pharmaceutically-acceptable salt thereof.

2. Compound of claim 1 wherein R^{28} is selected from (a) amino optionally substituted with two radicals selected from lower alkyl, lower aralkyl, lower heterocyclalkyl, heterocycl, and aryl, wherein R^{29} is selected from



wherein R^{30} is selected from lower alkyl, lower alkoxy, lower alkylamino, lower carboxyalkyl, lower alkoxyalkyl, lower alkylaminoalkyl, lower cycloalkyl, heterocyclyl, lower heterocyclalkyl, lower heterocyclalkoxy, lower aralkenyl, lower aralkyl, lower aralkoxy, phenyloxy, phenylamino, lower cycloalkyloxy, lower N-phenyl-N-alkylamino, lower alkylaminoalkoxy, lower alkylaminoalkylamino, lower heterocyclalkylamino, and lower N-aralkylamino; wherein R^{31} is lower alkyl; wherein R^{32} is selected from lower alkyl and aryl; and

wherein R^{33} is selected from hydrido and lower alkyl; or a pharmaceutically-acceptable salt thereof.

3, Compound of claim 2 selected from compounds and their pharmaceutically-acceptable salts, of the group consisting of

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[methyl(phenylmethyl)-amino]-4H-3,1-benzoxazin-4-one;

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[methyl[(4-methoxyphenyl)-methyl]amino]-4H-3,1-benzoxazin-4-one;

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[1-(1,2,3,6-tetrahydropyridyl)]-4H-3,1-benzoxazin-4-one;

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[isopropyl(methyl)amino]-4H-3,1-benzoxazin-4-one;

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[(4-furoyl)piperazinyl]-4H-3,1-benzoxazin-4-one;

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[[2-(2-cyano)ethyl]-(cyclopropyl)amino]-4H-3,1-benzoxazin-4-one;

6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[(4-thenoyl)piperazinyl]-4H-3,1-benzoxazin-4-one;


17 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[(4-benzenesulfonyl)-
18 piperazinyl]-4H-3,1-benzoxazin-4-one;
19 6-(benzoylamino)-5-methyl-2-[methyl(phenylmethyl)amino]-4H-3,1-benzoxazin-4-
20 one;
21 6-[(Phenylmethoxyacetyl)amino]-5-methyl-2-(diisopropylamino)-4H-3,1-
22 benzoxazin-4-one;
23 6-[(2,4,6-Trifluorobenzoyl)amino]-5-methyl-2(diisopropylamino)-4H-3,1-
24 benzoxazin-4-one;
25 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[[methyl(4-dimethylamino)-
26 phenylmethyl]-amino]-4H-3,1-benzoxazin-4-one;
27 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[methyl(2-pyridylmethyl)-
28 amino]-4H-3,1-benzoxazin-4-one;
29 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[methyl[2-(3-
30 indolyl)ethyl]amino]-4H-3,1-benzoxazin-4-one;
31 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-(4-morpholyl)-4H-3,1-
32 benzoxazin-4-one;
33 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[allyl(methyl)amino]-4H-3,1-
34 benzoxazin-4-one;
35 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-(1-decahydroquinolyl)-4H-
36 3,1-benzoxazin-4-one; and
37 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[4-(1-acetylpiperadiny)]-4H-
38 3,1-benzoxazin-4-one.

1 4. A pharmaceutical composition comprising a therapeutically-effective amount
2 of a compound of claim 1 and a pharmaceutically acceptable carrier or diluent.

1 5. A pharmaceutical composition comprising a therapeutically-effective amount
2 of a compound of claim 2 and a pharmaceutically acceptable carrier or diluent.




1 6. A pharmaceutical composition comprising a therapeutically-effective amount
2 of a compound of claim 3 and a pharmaceutically acceptable carrier or diluent.

7. A method or therapeutic or prophylactic treatment of Herpes Simplex Virus in a subject, said method comprising treating said subject with an effective amount of a compound of Formula II:




II

wherein R^{28} is selected from (a) amino optionally substituted with two radicals selected from alkyl, aralkyl, heterocyclalkyl, heterocycl, and aryl;
wherein R^{29} is selected from



wherein R^{30} is selected from alkyl, alkoxy, alkylamino, carboxyalkyl, alkoxyalkyl, alkylaminoalkyl, cycloalkyl, heterocycl, heterocyclalkyl, aralkyl, aralkoxy, aryloxy, cycloalkyloxy, arylamino, aralkenyl, heterocyclalkoxy, alkylaminoalkoxy, alkylaminoalkylamino, heterocyclalkylamino, N-aryl-N-alkylamino, and N-aralkylamino; wherein R^{31} is alkyl; wherein R^{32} is selected from alkyl and aryl; and wherein R^{33} is selected from hydrido, halo and alkyl;
or a pharmaceutically-acceptable salt thereof.

8. The method of 7 wherein R^{28} is selected from (a) amino optionally substituted with two radicals selected from lower alkyl, lower aralkyl, lower heterocyclalkyl, heterocycl, and aryl, wherein R^{29} is selected from



wherein R^{30} is selected from lower alkyl, lower alkoxy, lower alkylamino, lower carboxyalkyl, lower alkoxyalkyl, lower alkylaminoalkyl, lower cycloalkyl, heterocyclyl, lower heterocyclylalkyl, lower heterocyclylalkoxy, lower aralkenyl, lower aralkyl, lower aralkoxy, phenyloxy, phenylamino, lower cycloalkyloxy, lower N-phenyl-N-alkylamino, lower alkylaminoalkoxy, lower alkylaminoalkylamino, lower heterocyclylalkylamino, and lower N-aralkylamino; wherein R^{31} is lower alkyl; wherein R^{32} is selected from lower alkyl and aryl; and

wherein R^{33} is selected from hydrido and lower alkyl; or a pharmaceutically-acceptable salt thereof.

9. The method of claim 8 wherein the compound is selected from compounds and their pharmaceutically-acceptable salts, of the group consisting of
 - 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[methyl(phenylmethyl)-amino]-4H-3,1-benzoxazin-4-one;
 - 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[methyl[(4-methoxyphenyl)-methyl]amino]-4H-3,1-benzoxazin-4-one;
 - 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[1-(1,2,3,6-tetrahydro-pyridyl)]-4H-3,1-benzoxazin-4-one;
 - 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[isopropyl(methyl)amino]-4H-3,1-benzoxazin-4-one;
 - 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[(4-furoyl)piperazinyl]-4H-3,1-benzoxazin-4-one;
 - 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[[[(2-cyano)ethyl]-(cyclopropyl)amino]-4H-3,1-benzoxazin-4-one;

- 15 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[(4-thenoyl)piperazinyl]-4H-
16 3,1-benzoxazin-4-one;
- 17 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[(4-benzenesulfonyl)-
18 piperazinyl]-4H-3,1-benzoxazin-4-one;
- 19 6-(benzoylamino)-5-methyl-2-[methyl(phenylmethyl)amino]-4H-3,1-benzoxazin-4-
20 one;
- 21 6-[(Phenylmethoxyacetyl)amino]-5-methyl-2-(diisopropylamino)-4H-3,1-
22 benzoxazin-4-one;
- 23 6-[(2,4,6-Trifluorobenzoyl)amino]-5-methyl-2-(diisopropylamino)-4H-3,1-
24 benzoxazin-4-one;
- 25 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[[methyl(4-dimethylamino)-
26 phenylmethyl]-amino]-4H-3,1-benzoxazin-4-one;
- 27 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[methyl(2-pyridylmethyl)-
28 amino]-4H-3,1-benzoxazin-4-one;
- 29 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[methyl[2-(3-
30 indolyl)ethyl]amino]-4H-3,1-benzoxazin-4-one;
- 31 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-(4-morpholyl)-4H-3,1-
32 benzoxazin-4-one;
- 33 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[allyl(methyl)amino]-4H-3,1-
34 benzoxazin-4-one;
- 35 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-(1-decahydroquinolyl)-4H-
36 3,1-benzoxazin-4-one; and
- 37 6-[[[(1,1-Dimethylethoxy)carbonyl]amino]-5-methyl-2-[4-(1-acetylpiperadiny)]-4H-
38 3,1-benzoxazin-4-one.
- 39

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 January 2001 (18.01.2001)

PCT

(10) International Publication Number
WO 01/03697 A1

(51) International Patent Classification⁷: **A61K 31/536**,
C07D 265/18

(74) Agents: **WOLFFE, Susan, A. et al.**; Banner & Witcoff,
Ltd., Eleventh floor, 1001 G Street, N.W., Washington, DC
20001-4597 (US).

(21) International Application Number: **PCT/US00/18817**

(22) International Filing Date: **11 July 2000 (11.07.2000)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/142,956 **12 July 1999 (12.07.1999)** **US**

(71) Applicants (for all designated States except US): **G.D. SEARLE & CO.** [US/US]; Corporate Patent Department,
P.O. Box 5110, Chicago, IL 60680-5110 (US). **ASAHI CHEMICAL INDUSTRY CO., LTD.** [JP/JP]; 632-1, Mi-
fuku, Ohito-cho, Tagata-gun, Shizuoka-ken 410-2321 (JP).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KAWANISHI, Masashi** [JP/JP]; 632-1 Mifuku, Ohito-cho, Tagata-gun, Shizuoka-ken 410-2324 (JP). **TAKAHASHI, Wataru** [JP/JP]; 632-1, Mifuku, Ohito-cho, Tagata-gun, Shizuoka-ken 410-2324 (JP).

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **2-AMINO-BENZOXAZINONES FOR THE TREATMENT OF HERPES SIMPLEX VIRUS**

(57) Abstract: Antiviral agents and methods for treating Herpes Simplex Virus.

WO 01/03697 A1

JOINT DECLARATION FOR PATENT APPLICATION

As the below named inventors, we hereby declare that:

Our residence, post office address and citizenship are as stated below next to our names;

We believe we are the original, first and joint inventors of the subject matter which is claimed and for which a patent is sought on the invention entitled 2-AMINO-BENZOXAZINONES FOR THE TREATMENT OF HERPES SIMPLEX VIRUS, the specification of which

- ☐ is attached hereto.
- ☐ was filed on _____ as Application Serial Number _____ and was amended on _____ (if applicable).
- ☒ was filed under the Patent Cooperation Treaty (PCT) and accorded International Application No. PCT/US00/18817, filed July 11, 2000, and amended on _____ (if any).

We hereby state that we have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

We hereby acknowledge the duty to disclose information which is material to patentability in accordance with Title 37, Code of Federal Regulations, §1.56(a).

Prior Foreign Application(s)

We hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application(s) for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Country	Application No.	Date of Filing (day month year)	Date of Issue (day month year)	Priority Claimed Under 35 U.S.C. §119

Prior United States Provisional Application(s)

We hereby claim priority benefits under Title 35, United States Code, §119(e)(1) of any U.S. provisional application listed below:

U.S. Provisional Application No.	Date of Filing (day month year)	Priority Claimed Under 35 U.S.C. §119(e)(1)
60/142,956	12 July 1999	Yes

Prior United States Application(s)

We hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, we acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application Serial No.	Date of Filing (Day, Month, Year)	Status Patented, Pending, Abandoned

Power of Attorney

And we hereby appoint, both jointly and severally, as my attorneys, all Banner & Witcoff, Ltd. attorneys indicated therein under PTO Customer Number #22907, with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office.

All correspondence and telephone communications should be addressed to:

Banner & Witcoff, Ltd.
Customer Number: 22907 (WDC)

We hereby declare that all statements made herein of our own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

1-00
Signature Masashi Kawanishi Date 17 May 2002
Full Name of First Inventor KAWANISHI Masashi
Family Name First Given Name Second Given Name
Residence (city, state or country) JAPAN Citizenship JAPAN
Post Office Address 632-1, Mifuku, Ohito-Cho, Tagata-Gun, Shizuoka-Ken 410-2324, JAPAN JPX

2-00
Signature Wataru Takahashi Date 17 May 2002
Full Name of Second Inventor TAKAHASHI Wataru
Family Name First Given Name Second Given Name
Residence (city, state or country) JAPAN Citizenship JAPAN
Post Office Address 632-1, Mifuku, Ohito-Cho, Tagata-Gun, Shizuoka-Ken 410-2324, JAPAN JPX

Signature _____ Date _____
Full Name of Third Inventor _____
Family Name First Given Name Second Given Name
Residence (city, state or country) _____ Citizenship _____
Post Office Address _____

Signature _____ Date _____
Full Name of Fourth Inventor _____
Family Name First Given Name Second Given Name
Residence (city, state or country) _____ Citizenship _____
Post Office Address _____

☐ Additional Inventor Information Sheet(s) Attached